



TIDE Canada

Tide Glufosinate Ammonium 200SL Herbicide and Desiccant

By incorporating Tide Glufosinate Ammonium 200 SL into your crop management program, you can achieve broad-spectrum weed control and effective crop desiccation. This non-selective herbicide provides rapid action against a wide range of weeds, making it an essential tool for growers looking to optimize harvest efficiency and manage weed pressure effectively.



Why Choose Tide Glufosinate Ammonium 200SL?

- *Effective against tough annual and perennial weeds*
- *Compatible with LibertyLink® crop systems*
- *Flexible application rates for various weed pressures*
- *Non-volatile formulation for safer application*
- *Compatible with multiple tank mix partners*

Active Ingredient: Glufosinate Ammonium – 200 g/L
Chemistry Group: Group 10 – Glutamine Synthetase Inhibitor (*Non-Selective Contact Herbicide and Desiccant*)

Product Benefits

BROAD-SPECTRUM CONTROL

Effectively controls a wide range of annual and perennial broadleaf and grassy weeds, making it a versatile herbicide for various cropping systems.

FAST-ACTING CONTACT HERBICIDE

Provides rapid burn-down effects, with visible signs of weed desiccation appearing within hours to a few days after application.

NON-SELECTIVE HERBICIDAL ACTION

Targets all green plant tissues, ensuring effective control in agricultural settings.

MINIMAL SOIL RESIDUAL ACTIVITY

Deactivates upon contact with soil, reducing the risk of carryover and allowing for flexible crop rotation.

RESISTANCE MANAGEMENT TOOL

As a Group 10 herbicide, it provides an alternative mode of action to help manage herbicide-resistant weed populations.

Weeds Controlled

Annual Grasses:

- Barnyard grass
- Green foxtail
- Yellow foxtail
- Wild oats

Broadleaf Weeds:

- Redroot pigweed
- Lamb's-quarters
- Wild mustard
- Kochia

For a comprehensive list of weeds controlled, refer to the product label.

Application Guidelines

- **Water Volume:** Apply using a minimum of 10-24 gal/acre of water for ground applications to ensure thorough coverage.
- **Adjuvants:** When tank-mixing, always include the recommended adjuvants as specified on the tank mix partner's label.
- **Rainfast Period:** Avoid application if rainfall is expected within 4 hours of application to ensure optimal efficacy.

Environmental and Safety Precautions

- **Buffer Zones:** Maintain appropriate buffer zones to protect sensitive terrestrial and aquatic habitats.
- **Protective Equipment:** Wear protective clothing, including long-sleeved shirts, long pants, chemical-resistant gloves, and eye protection during mixing, loading, application, clean-up, and repair.
- **Re-Entry Interval:** Do not enter treated fields for 24 hours following application.

For detailed environmental precautions and first aid measures, consult the product label.

Packaging

- 2 x 10L jugs per carton
- 500L IBC totes
- 1000L IBC totes

REGISTERED CROPS AND APPLICATION RECOMMENDATIONS

CROP	APPLICATION RATE (L/ACRE)	APPLICATION TIMING	NOTES
Glufosinate-Ammonium Tolerant Field Corn	0.6 -0.8 L	Apply from the 1 to 8 leaf stage of corn.	Ensure thorough coverage for optimal weed control.
Glufosinate-Ammonium Tolerant Soybeans	0.6 -0.8 L	Apply from the cotyledon to flowering stage of soybeans.	Do not apply within 60 days of harvest.
Glufosinate-Ammonium Tolerant Canola (Eastern Canada and British Columbia)	0.6 -0.8 L	Apply from the cotyledon to early bolting stage of canola.	Do not apply within 65 days of harvest.

Maximum of two applications per year, with a minimum interval of 10 days between applications.

TANK MIX PARTNERS AND RATES

CROP	TANK MIX PARTNER	RATE OF TIDE GLUFOSINATE AMMONIUM 200SL (L/ACRE)	RATE OF TANK MIX PARTNER (L/ACRE)	NOTES
Glufosinate-Ammonium Tolerant Canola	TIDE Clethodim 240EC + TIDE Plant Guard or Centurion® or Select® + Amigo Adjuvant (0.5% v/v)	0.8 - 1.0	0.025	For control of Volunteer Barley and improved control of Wild Oats. Apply when weeds are at the 1-4 leaf stage with a maximum of 2 tillers.
Glufosinate-Ammonium Tolerant Corn	AAtrex® Liquid 480	0.8 - 1.0	0.71 - 0.94	Adjust atrazine rate to achieve desired residual control. This tank mix does not control later flushes of triazine-resistant weeds.
Glufosinate-Ammonium Tolerant Corn	Distinct®	0.8 - 1.0	0.115	Provides enhanced control of labeled annual broadleaf weeds.
Glufosinate-Ammonium Tolerant Corn	Marksman®	0.8 - 1.0	1.0 - 1.5	Provides residual control of annual grasses and annual broadleaf weeds. Adjust rate based on level of residual control desired.
Glufosinate-Ammonium Tolerant Soybeans	Pursuit®	0.8 - 1.0	0.125	Provides residual control of annual grasses and annual broadleaf weeds.
Glufosinate-Ammonium Tolerant Soybeans	Basagran® Forte	0.8 - 1.0	0.71	Provides enhanced control of annual broadleaf weeds.

Ensure compatibility and follow all label directions for both products when preparing tank mixtures.

Use of this product must comply with the label requirements. Always check and follow the most up-to-date label before application.



TIDE Canada

1-(866) 699-7717

info@tide-canada.com

www.tide-canada.com

Scott Jacques, Western Territory Account Manager
+1 (780) 690-1150 | scott.jacques@tide-canada.com

Jort Hopman, Country Manager
+1 (204) 384-9275 | jort.hopman@tide-canada.com